



**Gastroenterology Specialists**

22 Westfield Avenue  
Ansonia, CT 06401  
(203) 736-9919

Jeffrey Dreznick MD  
Sarah Olivier MD  
Jonathan Simon MD

**SUPREP COLONOSCOPY PREP**

This procedure will be performed at Griffin Hospital, 130 Division St., Derby, CT

**PLEASE BRING A CURRENT LIST OF MEDICATIONS**

Patient Name \_\_\_\_\_

**Report to the Outpatient Registration Desk on:**

Date \_\_\_\_\_ Time \_\_\_\_\_

- You must maintain a clear liquid diet all day starting the day before the procedure.
- Non-steroidal agents (Advil-Motrin-Ibuprofen) must be stopped 1 day prior to procedure.
- You are not allowed to drive. You are not allowed to go home via taxi by yourself.
- You must be accompanied by a friend or relative to assist you home.
- You may take your regular blood pressure/ heart/ thyroid/ seizure medication with a small sip of water at least 3 hours prior to your procedure unless instructed otherwise.
- Nothing to eat or drink three (3) hours before your procedure.
- If on blood thinners (Coumadin, Plavix, etc.) stop on \_\_\_\_\_.

Please bring these instructions to the G.I. Endoscopy Suite on the day of the procedure, along with a list of all your medications.

Please be aware that cancellation of this procedure will cause a delay in rescheduling.

**If you have any questions, please do not hesitate to call our office at (203) 736-9919**

## THE DAY BEFORE THE PROCEDURE:

- **Non-steroidal agents (Advil-Motrin-Ibuprofen) must be stopped 1 day prior to procedure.** It is safe to take Tylenol (acetaminophen) as a pain reliever. Consult with a physician on all other medications.
- Remain on a clear liquid diet. Solid foods, alcohol, milk or milk products are **not** allowed. Nothing red or purple in color. **It is very important to stay hydrated!**

### CLEAR LIQUIDS YOU MAY HAVE INCLUDE:

Black coffee (no milk or cream)

Tea (no milk or cream)

Chicken/beef/veggie broth

Gatorade/Powerade

Crystal Light

Water

Any type of soda

Snapple

Popsicles

Italian Ice

Jell-O

Clear fruit juices only – no orange juice

## COLON PREPARATION INSTRUCTIONS

- At \_\_\_\_\_ - Pour one (1) 6 ounce bottle of Suprep liquid into the mixing container. Add cool drinking water or **white cranberry juice** to the 16 ounce line on the container and mix well.
- Drink all of the liquid in the container and follow with two (2) more 16-ounce containers of clear liquids. Finish everything within one hour.
- If you have severe discomfort or distention (bloating), stop drinking the solution for a while or wait longer between drinking until the discomfort goes away.
- **Stay hydrated!** It's important that you drink clear liquids before, during and after your prep.

## THE DAY OF THE PROCEDURE:

- At \_\_\_\_\_ - Repeat the steps done the evening before using the other 6-ounce bottle.
- You must finish drinking the final glass of water at least three (3) hours before your procedure. After this is done you cannot eat or drink anything until the procedure is over.
- You may take your regular blood pressure/heart/thyroid/seizure medication with a small sip of water at least 3 hours prior to your procedure unless instructed otherwise.
- You **must** be accompanied by a friend or relative to drive and/or assist you home.

## TO RESCHEDULE:

As our patient we are committed to your health, and have reserved this procedure time exclusively for you. Please give us 48-hours notice if you wish to change this appointment. If you have any questions regarding the above instructions, please do not hesitate to contact our office at (203) 736-9919.